




Vaccine recommendations for the Abingdon VQA Feeder Cattle Sales

Components for <u>CALVES ON COWS</u>	Cows are not vaccinated or killed vaccines are used for respiratory viruses. (Choose 1 product for calves)	OR	Cow are vaccinated annually with modified live products. (Choose 1 product for calves)
<p><u>Round #1</u></p> <ol style="list-style-type: none"> 1. Clostridials (7-way blackleg) 2. Respiratory viruses (IBR, PI3, BRSV, BVD) 3. Dewormed (Pour-on or injectible) 	<p>TRIANGLE™ 5 <i>with Enhance™</i></p> <p>OR</p>	OR	<p>PYRAMID® 5</p> <p>OR</p>
<p><u>Could be given:</u></p> <ol style="list-style-type: none"> 1. At turn-out 2. Mid-summer 3. 2-weeks before weaning 4. Anytime in between! 		OR	

Vaccine recommendations for the Abingdon VQA Feeder Cattle Sales

Components for CALVES at WEANING	Modified-live Respiratory + M. Haemolytica (Choose 1 product for calves)
<ol style="list-style-type: none"> 1. Clostridials (7-way blackleg) 2. Respiratory viruses + Mannheimia haemolytica (IBR, PI3, BRSV, BVD) 3. Dewormed (Pour-on or Injectable) 	
<p style="text-align: center;"><u>Could be given:</u></p> <ol style="list-style-type: none"> 1. During weaning but at least 21 days prior to delivery. 	<p>or</p> 